



Assessment

Development and validation of the General Health Numeracy Test (GHNT)

Chandra Y. Osborn^{a,b,*}, Kenneth A. Wallston^c, Adam Shpigel^a, Kerri Cavanaugh^a, Sunil Kripalani^a, Russell L. Rothman^a

^a Department of Medicine, Vanderbilt University Medical Center, Nashville, USA

^b Department of Biomedical Informatics, Vanderbilt University Medical Center, Nashville, USA

^c School of Nursing, Vanderbilt University Medical Center, Nashville, USA

ARTICLE INFO

Article history:

Received 19 June 2012

Received in revised form 26 November 2012

Accepted 1 January 2013

Keywords:

General Health Numeracy Test

Validation

Measurement

Numeracy

Health literacy

Medication understanding

Medication adherence

ABSTRACT

Objective: Because existing numeracy measures may not optimally assess ‘health numeracy’, we developed and validated the General Health Numeracy Test (GHNT).

Methods: An iterative pilot testing process produced 21 GHNT items that were administered to 205 patients along with validated measures of health literacy, objective numeracy, subjective numeracy, and medication understanding and medication adherence. We assessed the GHNT’s internal consistency reliability, construct validity, and explored its predictive validity.

Results: On average, participants were 55.0 ± 13.8 years old, 64.9% female, 29.8% non-White, and 51.7% had incomes $\leq \$39$ K with 14.4 ± 2.9 years of education. Psychometric testing produced a 6-item version (GHNT-6). The GHNT-21 and GHNT-6 had acceptable-good internal consistency reliability (KR-20 = 0.87 vs. 0.77, respectively). Both versions were positively associated with income, education, health literacy, objective numeracy, and subjective numeracy (all $p < .001$). Furthermore, both versions were associated with participants’ understanding of their medications and medication adherence in unadjusted analyses, but only the GHNT-21 was associated with medication understanding in adjusted analyses.

Conclusions: The GHNT-21 and GHNT-6 are reliable and valid tools for assessing health numeracy.

Practice implications: Brief, reliable, and valid assessments of health numeracy can assess a patient’s numeracy status, and may ultimately help providers and educators tailor education to patients.

© 2013 Elsevier Ireland Ltd. All rights reserved.

1. Introduction

Literacy includes a host of skills including print literacy, oral literacy, and quantitative (numeracy) skills. Numeracy is an important component of literacy, and reflects one’s ability to understand and use numbers in daily life [1]. A growing body of evidence suggests individuals with limited literacy or numeracy skills are less likely to get preventive health care services [2,3], and are more likely to experience difficulty following medical instructions [4,5] understanding health information [6], performing self-care activities [7,8], and to have worse health outcomes [1,9] compared to individuals with adequate literacy or numeracy skills. For example, among individuals with diabetes, those with limited numeracy skills report worse diabetes knowledge and self-care activities, and have worse glycemic control compared to

individuals with adequate numeracy skills [10]. Finally, while literacy and numeracy are strongly correlated, some individuals may have adequate literacy but inadequate numeracy skills [4,6], and there are instances when numeracy, but not literacy, has been related to outcomes [10–12].

Most studies addressing literacy in health care use the term health literacy to refer to “one’s ability to understand and act on health information” [13] and include health literacy-specific measures, some of which are brief and could be administered in clinical care [14–16]. In contrast, some numeracy assessments used in research have limited application to health care decision making, or are lengthy mathematical tests that cannot be feasibly administered in clinical care [17,18]. Furthermore, there are a few health-related numeracy measures available, but these measures either assess a narrow range of mathematical skills [19,20] or are disease-specific [21,22], which limits our understanding of the cross-situational and cross-conditional impact of limited ‘health numeracy’ on health outcomes. Therefore, new measures to better estimate ‘health numeracy’ or “one’s understanding and capacity to act on numerical health information” are needed [1].

Provision of preventative, diagnostic, and therapeutic recommendations to patients often includes understanding and using

* Corresponding author at: Vanderbilt University Medical Center, Division of General Internal Medicine & Public Health, Center for Health Services Research, 1215 Twenty-First Avenue South, Suite 6000, MCE – North Tower, Nashville, TN 37232-8300, USA. Tel.: +1 615 936 8468; fax: +1 615 936 1269.

E-mail address: chandra.osborn@vanderbilt.edu (C.Y. Osborn).

applied quantitative skills such as executing a medication schedule, or dietary recommendations and understanding results of laboratory assessments. In addition, numeracy is fundamental for the delivery of patient-centered care where options in management, including potential risks and benefits of these choices may be discussed [23]. Identification of patient's with limited health numeracy skills may be important to trigger targeted resources to ensure a valuable experience for both patients and providers. Of particular interest would be tools with application in a variety of health care contexts that are brief, well-received by patients, and have practical, clinical and scientific utility for providers and researchers.

Our study objective was to develop an assessment of health numeracy, referred to as the General Health Numeracy Test (GHNT) and evaluate its psychometric properties, including internal consistency reliability and construct validity. Given recent evidence suggesting numeracy skills are strongly associated with one's capacity to manage medications, and even explain gender and racial disparities in medication management [24,25], we also explored the GHNT's predictive validity with patient's understanding of their medications and medication adherence.

2. Methods

2.1. Sample

From June 2009 to August 2010, trained research assistants (RAs) enrolled 205 patients from the Vanderbilt Adult Primary Care Clinic in Nashville, TN. Eligible patients were between the ages of 18 and 80 years and English-speaking. RAs used the Rosenbaum Pocket screener [26] to assess visual acuity, and excluded patients whose visual acuity was worse than 20/50. Patients were also excluded if they were too ill to participate, or had a clinical diagnosis of blindness, dementia and/or psychosis.

2.2. Data and procedure

Trained RAs recruited eligible patients, consented patients who were interested in participation, and conducted individual, in-person interviews to collect demographic information, administer validated measures of health literacy, objective numeracy, subjective numeracy, medication understanding, medication adherence, and the newly developed GHNT. In an effort to minimize the impact of patients' health literacy status on their numeracy skills, RAs first verbally administered the Rapid Estimate of Adult Learning in Medicine (REALM). If a patient scored ≤ 44 on the REALM (i.e., they were categorized as having low health literacy), the remaining measures were verbally administered. Otherwise, RAs offered patients the option of verbal administration or self-administration. All interviews were conducted in a private room immediately prior to or after a patient's scheduled medical appointment.

2.2.1. Demographics

Self-reported socio-demographic information included patient age, gender, race, income, years of education, and health insurance status.

2.2.2. Health literacy

The REALM is a valid and reliable measure of health literacy [27–29]. Respondents are asked to read aloud 66 health-related words arranged in order of increasing difficulty [27]. The score assigns health literacy skills into four categories of grade-equivalent reading level. Correct pronunciation of 0–44 words represents a 6th grade reading level or below (low literacy), 45–60 words represents a 7th–8th grade reading level (marginal literacy),

and 61–66 words represents a 9th grade reading level (adequate literacy).

2.2.3. Objective numeracy

The arithmetic section of the Wide Range Achievement Test, 3rd Edition (WRAT-3) is a validated instrument that evaluates the ability to compute solutions to math problems [17]. Part I requires counting, reading number symbols, and solving simple arithmetic problems that are verbally presented to the examinee. Part II consists of using paper and a pencil to calculate up to 40 arithmetic problems within 15 min. Points (1 for each correct answer) are summed across Parts I and II, combined into a single raw score, and then standardized by age.

2.2.4. Subjective numeracy

The Subjective Numeracy Scale (SNS) is a valid and reliable self-report measure of one's perceived ability to perform various mathematical tasks and one's preference for using numerical versus prose information [20,30]. This 8-item scale requires no mathematical calculations. Instead, it consists of 4 items asking respondents to assess their numerical ability in different contexts and 4 items asking respondents to state their preference for the presentation of numerical and probabilistic information.

2.2.5. Medication understanding

The Medication Understanding Questionnaire (MUQ) is a valid and reliable measure of one's knowledge of the purpose, dose, and frequency of the medication(s) in his regimen [31]. To administer the MUQ, RAs access a patient's medication list from the medical record and use a random numbers table to select up to 5 prescription medications from this list. If a patient is taking ≤ 5 medications, all eligible medications are selected. In our study, eligible medications were those taken every day, excluding medications that were taken 'as needed,' were considered combination therapies, or were classified as lotions/creams, nasal sprays, eye drops, herbals, supplements, or vitamins. The RA provides the brand and generic name of medication, and then asks the patient for the medication's purpose, strength per unit (e.g., 20 mg tablet), number of units taken at a time (e.g., 2 tablets), and dosing frequency (e.g., twice a day). Patients are instructed not to refer to their medication list or bottles when responding. Each medication is scored from 0 to 3, reflecting the medication's indication (1 point), strength (0.5 point), units (0.5 point), and frequency (1 point). While a patient's overall MUQ score is the average of the scores for each tested medication, we converted these scores to percent correct to be consistent with our measure of medication adherence.

2.2.6. Medication adherence

The Summary of Diabetes Self-Care Activities (SDSCA) medications subscales (i.e., the 1 item and 2 item versions) are valid and reliable measures of self-reported medication adherence, and individual items have correlated with objective measures of medication adherence (e.g., medication possession ratio) [32]. We used the single item version modified to make it applicable to patients without diabetes and any drug indication. In the current study, we presented patients with each medication in their regimen for up to 5 randomly selected, eligible medications. If a patient is taking ≤ 5 medications, all eligible medications are selected. In our study, eligible medications were those taken every day, excluding medications that were taken 'as needed,' were considered combination therapies, or were classified as lotions/creams, nasal sprays, eye drops, herbals, supplements, or vitamins. We then asked the patient, "Please tell me how many days in the last week you took your (insert one medication name)." We then calculated each patient's percent adherence score (i.e., [(number of

medications queried up to 5) (7 days) – (total days any medication was missed)/[(number of medications queried up to 5) (7 days)] = % adherence) [33]. This was treated as a non-parametric continuous variable [32,34] and also categorized as adherent (100%) or not (<100%) [32,35,36].

2.2.7. General Health Numeracy Test

Six experts in health literacy, numeracy, health psychology, scale development, and clinical medicine developed a novel set of 63 health numeracy items after reviewing previously developed assessment tools. These items assessed a wide range of numerical skills often used in the context of making health decisions; for example, understanding number hierarchy, performing calculations and correctly estimating probability when making decisions about disease prevention, nutrition management, medication adherence, and assessing one's risk of experiencing a negative health outcome. The expert panel reviewed all items, omitting those items assessing the same numerical skill with the same level of difficulty. This resulted in a set of 44 items that were divided into 4 forms with 11 items per form. We administered each form to 6 patients ($n = 24$), and performed cognitive interviews with these patients to further identify those items assessing the same numerical skill with the same level of difficulty. Findings allowed us to further reduce the set to 33 items. We then performed a second round of testing, administering all 33 items to a sample of patients ($n = 35$) and using a combination of cognitive interviews with these patients and expert panel review to further reduce the set to 21 items (GHNT-21). During each iterative, pilot testing phase, we omitted items that were unclear to patients, too difficult to answer, or were redundant with other items, while purposefully retaining items that covered a broad range of numerical skills described above.

2.3. Data analyses

All analyses were performed using SPSS version 21.0. In an effort to further reduce the number of items on the GHNT-21, we first performed multiple rounds of psychometric testing (i.e., calculating internal consistency reliabilities if items were deleted and conducting tests of construct validity and predictive validity). These analyses resulted in a 6-item version of the GHNT (GHNT-6) shown in Fig. 1.

The Kuder–Richardson (KR-20) coefficient was used to examine the GHNT-21's and GHNT-6's internal consistency reliabilities. Values range from 0 to 1, with a higher KR-20 (e.g., >0.90) indicating a homogenous test.

A series of Spearman's rho correlation coefficients tested our a priori hypothetical model for construct validity. This model predicted that health numeracy measured with the GHNT-21 or GHNT-6 would be positively associated with patients' income, years of education, health literacy, objective numeracy, and subjective numeracy.

To explore the GHNT-21's and GHNT-6's predictive validity, we postulated that higher GHNT scores would correspond with greater medication understanding and medication adherence. To test these hypothesis, we performed a series of multivariate linear regression models to test the unadjusted and adjusted relationships between the GHNT-21 or the GHNT-6 and medication understanding and medication adherence as continuous variables. We then performed multivariate logistic regression models to test the unadjusted and adjusted relationship between the GHNT-21 or the GHNT-6 and medication adherence as a dichotomized variable (i.e., coded as 0 = perfect adherence and 1 = less than perfect adherence). In all adjusted models, we adjusted for age, gender, racial status, and income. We did not adjust for education,

insurance status, or employment status due to these variables being collinear with numeracy.

3. Results

3.1. Sample characteristics

Sample characteristics are presented in Table 1. On average, participants ($N = 205$) were 55 years old ($SD = 13.8$ years), 64.9% were female, 29.8% were non-White, and 69.8% reported having private health insurance. Sixty-seven percent reported at least some college education, while 51.7% reported a total annual household income ≤ 39 K. Eighteen percent of the sample had <9th grade reading level (or limited health literacy skills) according to the REALM, but 61.8% of the sample had <9th grade numeracy skills according to the WRAT-3. Finally, on average, participants understood their medication regimen 81.7% of the time and were adherent to their regimen 90.7% of the time (i.e., medication adherence as a continuous variable), with 67.5% of the sample reporting perfect adherence (i.e., medication adherence as a categorical variable, 100% adherence vs. <100% adherence).

3.2. Descriptive statistics, reliability, and construct validity

The mean percent correct score on the GHNT-21 was 56% (out of 100%) with a standard deviation of 23%, which was slightly higher than the neutral midpoint (50%). The distribution was near normal, with only a slight negative skew (skewness -0.19 , SE of skewness 0.17) and slight platykurtic spread (kurtosis -0.74 , SE of kurtosis 0.34). Internal consistency reliability was good according to the Kuder–Richardson coefficient (KR-20 = 0.87). As shown in Table 2, higher scores on the GHNT-21 were significantly associated with higher income, more years of education, higher health literacy scores, higher objective numeracy scores, and higher subjective numeracy scores ($p < .001$ for all comparisons).

The mean percent correct score on the GHNT-6 (see Fig. 1 for percent correct per item) was 42% (out of 100%) with a standard deviation of 30%, which was slightly lower than the neutral midpoint (50%). The distribution was near normal, with only a slight negative skew (skewness 0.43, SE of skewness 0.17) and slight platykurtic spread (kurtosis -1.01 , SE of kurtosis 0.34). Internal consistency reliability was acceptable according to the Kuder–Richardson coefficient (KR-20 = 0.77). As shown in Table 2, higher scores on the 6-item GHNT were significantly associated with higher income, more years of education, higher health literacy scores, higher objective numeracy scores, and higher subjective numeracy scores ($p < .001$ for all comparisons).

3.3. Predictive validity

As shown in Table 3, higher scores on the GHNT-21 were significantly associated with medication understanding in both the unadjusted, $F = 28.55$, $p < .001$ ($\beta = 0.36$, $p < .001$), and adjusted models, $F = 7.99$, $p < .001$ ($\beta = 0.29$, $p < .001$). In unadjusted models, higher scores on the GHNT-21 were significantly associated with both medication adherence as a continuous variable, $F = 5.42$, $p = .02$ ($\beta = 0.17$, $p = .02$), and as a dichotomized variable (adherent vs. non-adherent; OR = 4.75, $p = .02$ [95% CI: 1.28–17.69]). However, in adjusted models, scores on the GHNT-21 were not associated with medication adherence as a continuous variable, $F = 3.37$, $p < .01$ ($\beta = 0.13$, $p = .18$), or as a dichotomized variable (adherent vs. non-adherent; AOR = 1.82, $p = .50$ [95% CI: 0.32–10.36]).

1. Call your doctor if you have a temperature of 100.4 °F or greater. The thermometer looks like the following:

100.2 F

Do you call the doctor?

ANSWER: _____ YES _____ NO [87.3% responded correctly]

2. If 4 people out of 20 have a chance of getting a cold, what would be the risk of getting a cold?

ANSWER: _____ % [36.1% responded correctly]

3. Suppose that the maximum heart rate for a 60 year old woman is 160 beats per minute and that she is told to exercise at 80% of her maximum heart rate. What is 80% of that woman's maximum heart rate?

ANSWER: _____ beats per minute [46.8% responded correctly]

4. You ate half the container of carrots. How many grams of carbohydrates did you eat?

Nutrition Facts	
Serving Size: 1 cup (85g) (3 oz.)	
Servings Per Container: 2.5	
Amount Per Serving	
Calories 45	Calories from Fat 0
% Daily Value*	
Total Fat 0g	0%
Saturated Fat 0g	0%
Cholesterol 0mg	0%
Sodium 55 mg	2%
Total Carbohydrate 10g	3%
Dietary Fiber 3g	12%
Sugars 5g	
Protein 1g	

ANSWER: _____ grams [26.3% responded correctly]

5. Your doctor tells you that you have high cholesterol. He informs you that you have a 10% risk of having a heart attack in the next 5 years. If you start on a cholesterol-lowering drug, you can reduce your risk by 30%.

What is your 5-year risk if you take the drug?

ANSWER: _____ % [21.0% responded correctly]

6. A mammogram is used to screen women for breast cancer. False positives are tests that incorrectly show a positive result. 85% of positive mammograms are actually false positives. If 1,000 women receive mammograms, and 200 are told there is an abnormal finding, how many women are likely to actually have breast cancer?

ANSWER: _____ women [35.1% responded correctly]

Fig. 1. The 6-item General Health Numeracy Test (GHNT-6).

As shown in Table 3, higher scores on the GHNT-6 were significantly associated with medication understanding in both the unadjusted, $F = 12.15$, $p < .001$ ($\beta = 0.25$, $p < .001$), and adjusted models, $F = 6.43$, $p < .001$ ($\beta = 0.16$, $p = .05$). In unadjusted models, higher scores on the GHNT-6 were marginally associated with medication adherence as a continuous variable, $F = 3.58$, $p = .06$ ($\beta = 0.14$, $p = .06$), and as a dichotomized variable (adherent vs. non-adherent; OR = 2.54, $p = .08$ [95% CI: 0.90–7.18]). However, in adjusted models, scores on the GHNT-6 were not associated with medication adherence as a continuous variable, $F = 3.14$, $p < .01$

($\beta = 0.07$, $p = .38$), or as a dichotomized variable (adherent vs. non-adherent; AOR = 1.23, $p = .74$ [95% CI: 0.36–4.27]).

4. Discussion and conclusion

4.1. Discussion

The GHNT-21 and GHNT-6 are reliable and valid measures of health numeracy. The GHNT-21 demonstrated better internal consistency reliability than the GHNT-6, but both measures were

Table 1
Participant characteristics.

N = 205	M ± SD or n (%)
Age	55.0 ± 13.8
Gender	
Female	133 (64.9)
Race	
Caucasian/White	144 (70.2)
African American/Black	53 (25.8)
Other	7 (4.0)
Education (years)	14.4 ± 2.9
<High school	13 (6.4)
High school	54 (26.6)
>High school	136 (67.0)
Income	
≤39K	104 (51.7)
≥40K	97 (48.3)
Insurance status	
Public/none	60 (29.3)
Private	143 (69.8)
Medical condition, range 0–3	1.1 ± 0.9
Diabetes mellitus	60 (29.4)
Hypertension	133 (65.2)
Heart disease	43 (21.1)
Health Literacy (REALM)	
<9th grade	37 (18.0)
≥9th grade	168 (82.0)
Objective Numeracy (WRAT-3)	
<9th grade	126 (61.8)
≥9th grade	78 (38.2)
Subjective Numeracy (SNS), range 1–6	3.8 ± 1.2
Medication Understanding (MUQ), % correct	81.7 ± 17.4
Medication Adherence (SDSCA), % adherent	90.7 ± 20.5
Perfectly adherent	133 (67.5)
Not perfectly adherent	64 (32.5)

Notes: REALM, Rapid Estimate of Adult Learning in Medicine; WRAT-3, Wide Range Achievement Test, 3rd Edition; SNS, Subjective Numeracy Scale; MUQ, Medication Understanding Questionnaire; SDSCA, Summary of Diabetes Self-Care Activities.

significantly associated with measures used to test for construct validity. While both measures were also associated with medication understanding and medication adherence, the 21-item version had better predictive validity with medication understanding and adherence than the 6-item version did. However, the 6-item version may require less time to administer and create less response burden, and thus may be an appropriate tool for identifying patients with limited health numeracy.

Our sample reported high rates of limited numeracy skills. While <20% of the sample had limited health literacy skills according to the REALM, approximately two thirds had limited numeracy skills according to the WRAT-3, and the average percent correct on the GHNT-21 and GHNT-6 were 56% and 42% (both out of 100%), respectively. On the GHNT-6, participants were more

likely to answer correctly to questions requiring fewer calculations or an understanding of number hierarchy, and were least likely to respond correctly to questions requiring more calculations or an understanding of risk and probability. For example, only 26.3% of the sample could correctly calculate the number of carbohydrates on the Nutrition Facts label, and only 21.0% could correctly calculate the risk of having a heart attack in 5 years if taking cholesterol-lowering medications that reduces this risk by a certain amount.

In adjusted analyses, the GHNT-21 was associated with patients understanding their medication regimen. Successful adherence to medications not only relies on the frequency of medication-taking across time, but also on accurate understanding and management of prescription instructions. This includes correct identification of pills, understanding the number of pills to be taken at each dose, the timing of doses, determination of missed doses, need for refills and comprehension of warnings and other ancillary instructions (e.g., “take on an empty stomach”) [37]. Widely used measures of health literacy have been consistently associated with both understanding one’s medication regimen [7,31,38] and medication adherence [16,39–42]. However, no studies to our knowledge have explored the relationship between numeracy and patients’ understanding of their own medication regimen, and only one study has found support for a relationship between objective numeracy measured with a non-health-related assessment and medication management capacity [24,25].

We recently tested the relationship between disease-specific numeracy (measured with the Diabetes Numeracy Test [21]) and medication adherence, and objective numeracy (measured with the WRAT-3 [18]) and medication adherence, and found no associations [39]. In contrast, the current study explored the relationship between numeracy measured with the new GHNT-21 or GHNT-6 and medication understanding and medication adherence, and found support for ‘health numeracy’ being associated with medication understanding and weaker support for ‘health numeracy’ being associated with medication adherence, with both providing initial evidence of the GHNT’s predictive validity. Future research should continue to explore these relationships using multiple measures of medication understanding and adherence, and larger samples that are homogenous with respect to medical condition and drug indication.

There are study limitations to acknowledge. First, we assessed medication adherence via a single, self-report measure rather than more objective measures of adherence (e.g., pill counts, prescription refills, and electronic monitoring devices). While patients may under-report missed doses through questionnaires, a handful of studies suggest self-report measures are viable and accurate measures of medication adherence [43,44].

Table 2
Inter-correlations of study variables.

Variables	GHNT-21	GHNT-6	Education	Income	REALM	WRAT-3	SNS	MUQ	SDSCA
GHNT-21	1.00								
GHNT-6	0.88***	1.00							
Education	0.45***	0.41***	1.00						
Income	0.34***	0.32***	0.39***	1.00					
REALM	0.54***	0.45***	0.42***	0.31***	1.00				
WRAT-3	0.64***	0.65***	0.45***	0.23***	0.47***	1.00			
SNS	0.57***	0.58***	0.52***	0.33***	0.36***	0.60***	1.00		
MUQ	0.35***	0.25***	0.28***	0.15*	0.36***	0.24***	0.08	1.00	
SDSCA	0.18*	0.16*	0.04	0.08	0.12	0.14*	0.07	0.22**	1.00

Notes: Spearman’s rho assessed the inter-correlations of study variables located in the remaining cells. GHNT-21, 21-item General Health Numeracy Test; GHNT-6, 6-item General Health Numeracy Test; REALM, Rapid Estimate of Adult Learning in Medicine; WRAT-3, Wide Range Achievement Test, 3rd Edition; SNS, Subjective Numeracy Scale; MUQ, Medication Understanding Questionnaire; SDSCA, Summary of Diabetes Self-Care Activities (medications subscale).

* $p < .05$.
** $p < .01$.
*** $p < .001$.

Table 3

Multivariate linear regression models with the 21-item and 6-item General Health Numeracy Tests (GHNT-21 and GHNT-6, respectively) predicting medication understanding.

Predictor	GHNT-21			GHNT-6				
	Unstandardized coefficient		β	<i>p</i> -Value	Unstandardized coefficient		β	<i>p</i> -Value
	<i>B</i>	SE			<i>B</i>	SE		
Model 1: dependent variable = medication understanding GHNT	26.78	5.01	0.36	.000	14.10	4.04	0.25	.001
Model 2: dependent variable = medication understanding GHNT	21.16	6.52	0.29	.001	8.98	4.63	0.16	.054
Age	−0.22	0.09	−0.17	.015	−0.28	0.09	−0.22	.001
Gender (male vs. female)	4.84	2.47	0.13	.052	5.44	2.53	0.15	.033
Race (Caucasian/White vs. African American/Black)	−2.28	3.23	−0.06	.481	−5.42	3.06	−0.14	.078
Income (≤ 39 K vs. ≥ 40 K)	1.12	2.58	0.03	.664	2.67	2.56	0.08	.299

Notes: GHNT, General Health Numeracy Test; B, unstandardized coefficient; SE, standard error; β , standardized coefficient.

Secondly, we do not know the administration time of the GHNT-6 since we administered the GHNT-21 in our main study. However, based on our experience administering numeracy instruments, we estimate this time to be approximately 5–8 min. The GHNT also focuses on computational skills and “verbatim” responses rather than on patients’ “number sense.” Reyna [45] has suggested that many patients understand numbers and make mathematical inferences based more on “gist” or “number sense” than on exact computations (or what Reyna refers to as “verbatim” representations of information). While there is often a correlation between a patient’s computational skill (and “verbatim” responses) and their “number sense,” there may be patients with good number sense, but poor computational skills and patients with good computational skills, but poor number sense. In addition, whether or not a patient can perform a set of computations to 100% accuracy may not be relevant to the task being tested or the communication being interpreted—as long as they have understood the “gist” of the situation and make appropriate actions in response.

Finally, our study sample was also small, predominately Caucasian/White race, of higher socioeconomic status and from a single academic medical center. Thus, the findings may not generalize to lower income, racially/ethnically diverse patient populations and/or those receiving care at non-academic facilities. Furthermore, as primary care patients, this sample was heterogeneous with respect to the number and type of medical conditions they had, the number of medications they were taking, and the indication of these drugs, potentially limiting our ability to detect a statistically significant association between the GHNT-6 and medication understanding and medication adherence in our adjusted analyses. Despite these limitations, our study is the first to report the psychometric properties and predictive validity of a measure of ‘health numeracy’ that represents a range of health topics.

4.2. Conclusion

While our findings suggest the GHNT-21 and GHNT-6 are valid and reliable measures of ‘health numeracy’, additional predictive validity evidence is needed to ascertain if the 6-item version is as robust as the 21-item version for research purposes. In general, future research should provide additional psychometric support as well as evidence of the utility of both instruments in intervention research and clinical care. Such evaluation might include establishing test–retest reliability, discriminant validity, scoring cut-offs to categorize individuals with limited versus adequate health numeracy, and providing evidence of the GHNT-21’s and GHNT-6’s predictive validity with a wide range of social-cognitive, behavioral, and physiological health outcomes. Finally, additional support for the instruments’ association with medication

understanding and medication adherence along with evidence of the instruments’ sensitivity would allow for determining whether the GHNT is an applicable evaluative tool for interventions promoting these outcomes.

4.3. Practice implications

Our findings demonstrate that a relatively brief tool can assess a patient’s health numeracy status and highlight the importance of understanding how a patient’s numeracy might impact their understanding of issues related to their health care and self-management, such as understanding a medication regimen and adhering to it. Recent studies suggest there may be opportunities to improve how providers communicate to patients with low health numeracy about taking care of their health, especially about self-management behaviors self-care activities requiring numeracy skills (e.g., correct timing and dosing of medications) [46,47]. However, in order to be effective, we must first develop, test, and incorporate brief, reliable, and valid assessments of health numeracy [1]. Such measures may ultimately help health care providers and educators tailor educational messages to a patients’ level of understanding, with the goal of improving patient self-care and health outcomes.

Conflict of interest

Dr. Kripalani is a consultant to and holds equity in PictureRx, LLC, which makes patient education tools to improve medication management. PictureRx did not provide materials or funding for this study.

Acknowledgments

Dr. Osborn is supported by NIDDK K01DK087894. Dr. Cavanaugh is supported by NIDDK K23DK80952.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.pec.2013.01.001>.

References

- [1] Rothman RL, Montori VM, Cherrington A, Pignone MP. Perspective: the role of numeracy in health care. *J Health Commun* 2008;13:583–95.
- [2] Ciampa PJ, Osborn CY, Peterson NB, Rothman RL. Patient numeracy, perceptions of provider communication, and colorectal cancer screening utilization. *J Health Commun* 2010;15:157–68.

- [3] Sudore RL, Mehta KM, Simonsick EM, Harris TB, Newman AB, Satterfield S, et al. Limited literacy in older people and disparities in health and healthcare access. *J Am Geriatr Soc* 2006;54:770–6.
- [4] Kumar D, Sanders L, Perrin EM, Lokker N, Patterson B, Gunn V, et al. Parental understanding of infant health information: health literacy, numeracy, and the Parental Health Literacy Activities Test (PHLAT). *Acad Pediatr* 2010;10:309–16.
- [5] Lindquist LA, Go L, Fleisher J, Jain N, Friesema E, Baker DW. Relationship of health literacy to intentional and unintentional non-adherence of hospital discharge medications. *J Gen Intern Med* 2012;27:173–8.
- [6] Rothman RL, Housam R, Weiss H, Davis D, Gregory R, Gebretsadik T, et al. Patient understanding of food labels: the role of literacy and numeracy. *Am J Prev Med* 2006;31:391–8.
- [7] Kripalani S, Henderson LE, Chiu EY, Robertson R, Kolm P, Jacobson TA. Predictors of medication self-management skill in a low-literacy population. *J Gen Intern Med* 2006;21:852–6.
- [8] Huizinga MM, Carlisle AJ, Cavanaugh KL, Davis DL, Gregory RP, Schlundt DG, et al. Literacy, numeracy, and portion-size estimation skills. *Am J Prev Med* 2009;36:324–8.
- [9] Wolf MS, Gazmararian JA, Baker DW. Health literacy and functional health status among older adults. *Arch Intern Med* 2005;165:1946–52.
- [10] Cavanaugh K, Huizinga MM, Wallston KA, Gebretsadik T, Shintani A, Davis D, et al. Association of numeracy and diabetes control. *Ann Intern Med* 2008;148:737–46.
- [11] Huizinga MM, Beech BM, Cavanaugh KL, Elasy TA, Rothman RL. Low numeracy skills are associated with higher BMI. *Obesity (Silver Spring)* 2008;16:1966–8.
- [12] Lokker N, Sanders L, Perrin EM, Kumar D, Finkle J, Franco V, et al. Parental misinterpretations of over-the-counter pediatric cough and cold medication labels. *Pediatrics* 2009;123:1464–71.
- [13] IOM. In: Nielsen-Bohlman L, Panzer A, Kindig DA, editors. *Health literacy: a prescription to end confusion*. Washington, DC: The National Academies Press; 2004.
- [14] Chew LD, Bradley KA, Boyko EJ. Brief questions to identify patients with inadequate health literacy. *Fam Med* 2004;36:588–94.
- [15] Morris NS, MacLean CD, Chew LD, Littenberg B. The single item literacy screener: evaluation of a brief instrument to identify limited reading ability. *BMC Fam Pract* 2006;7:21.
- [16] Osborn CY, Davis TC, Bailey SC, Wolf MS. Health literacy in the context of HIV treatment: introducing the brief estimate of health knowledge and action (BEHKA)-HIV version. *AIDS Behav* 2010;14:181–8.
- [17] Wilkinson GS. *WRAT3: word range achievement test administration manual*. Wilmington, DE: Wide Range Inc.; 1993.
- [18] Woodcock R, McGrew K, Mather N. *Woodcock Johnson III tests of achievement*. Itasca, IL: Riverside Publishing; 2001.
- [19] Lipkus IM, Samsa G, Rimer BK. General performance on a numeracy scale among highly educated samples. *Med Decis Making* 2001;21:37–44.
- [20] Zikmund-Fisher BJ, Smith DM, Ubel PA, Fagerlin A. Validation of the Subjective Numeracy Scale: effects of low numeracy on comprehension of risk communications and utility elicitations. *Med Decis Making* 2007;27:663–71.
- [21] Huizinga MM, Elasy TA, Wallston KA, Cavanaugh K, Davis D, Gregory RP, et al. Development and validation of the Diabetes Numeracy Test (DNT). *BMC Health Serv Res* 2008;8:96.
- [22] Apter AJ, Cheng J, Small D, Bennett IM, Albert C, Fein DG, et al. Asthma numeracy skill and health literacy. *J Asthma* 2006;43:705–10.
- [23] Committee on Quality of Health Care in America. *Crossing the quality chasm: a new health system for the 21st century*. Washington, DC: Institute of Medicine; 2001.
- [24] Waldrop-Valverde D, Jones DL, Jayaweera D, Gonzalez P, Romero J, Ownby RL. Gender differences in medication management capacity in HIV infection: the role of health literacy and numeracy. *AIDS Behav* 2009;13:46–52.
- [25] Waldrop-Valverde D, Osborn CY, Rodriguez A, Rothman RL, Kumar M, Jones DL. Numeracy skills explain racial differences in HIV medication management. *AIDS Behav* 2010;14:799–806.
- [26] Rosenbaum J. *Rosenbaum vision pocket screener*. Quincy, MA: Grass Instruments; 1982.
- [27] Davis TC, Long SW, Jackson RH, Mayeaux EJ, George RB, Murphy PW, et al. Rapid estimate of adult literacy in medicine: a shortened screening instrument. *Fam Med* 1993;25:391–5.
- [28] Parker RM, Baker DW, Williams MV, Nurss JR. The test of functional health literacy in adults: a new instrument for measuring patients' literacy skills. *J Gen Intern Med* 1995;10:537–41.
- [29] Davis TC, Michielutte R, Askov EN, Williams MV, Weiss BD. Practical assessment of adult literacy in health care. *Health Educ Behav* 1998;25:613–24.
- [30] Fagerlin A, Zikmund-Fisher BJ, Ubel PA, Jankovic A, Derry HA, Smith DM. Measuring numeracy without a math test: development of the Subjective Numeracy Scale. *Med Decis Making* 2007;27:672–80.
- [31] Marvanova M, Roumie CL, Eden SK, Cawthon C, Schnipper JL, Kripalani S. Health literacy and medication understanding among hospitalized adults. *J Hosp Med* 2011;6:487.
- [32] Cohen HW, Shmukler C, Ullman R, Rivera CM, Walker EA. Measurements of medication adherence in diabetic patients with poorly controlled HbA(1c). *Diabet Med* 2010;27:210–6.
- [33] Cohen MJ, Shaykevich S, Cawthon C, Kripalani S, Paasche-Orlow MK, Schnipper JL. Predictors of medication adherence postdischarge: the impact of patient age, insurance status, and prior adherence. *J Hosp Med* 2012;7:470–5.
- [34] Toobert DJ, Hampson SE, Glasgow RE. The summary of diabetes self-care activities measure: results from 7 studies and a revised scale. *Diabetes Care* 2000;23:943–50.
- [35] Gonzalez JS, Safren SA, Cagliero E, Wexler DJ, Delahanty L, Wittenberg E, et al. Depression, self-care, and medication adherence in type 2 diabetes: relationships across the full range of symptom severity. *Diabetes Care* 2007;30:2222–7.
- [36] Gonzalez JS, Safren SA, Delahanty LM, Cagliero E, Wexler DJ, Meigs JB, et al. Symptoms of depression prospectively predict poorer self-care in patients with Type 2 diabetes. *Diabet Med* 2008;25:1102–7.
- [37] MacLaughlin EJ, Raehl CL, Treadway AK, Sterling TL, Zoller DP, Bond CA. Assessing medication adherence in the elderly: which tools to use in clinical practice? *Drugs Aging* 2005;22:231–55.
- [38] Wolf MS, Davis TC, Shrank W, Rapp DN, Bass PF, Connor UM, et al. To err is human: patient misinterpretations of prescription drug label instructions. *Patient Educ Couns* 2007;67:293–300.
- [39] Osborn CY, Cavanaugh K, Wallston KA, Kripalani S, Elasy TA, Rothman RL, et al. Health literacy explains racial disparities in diabetes medication adherence. *J Health Commun* 2011;16:268–78.
- [40] Kalichman SC, Pope H, White D, Cherry C, Amaral CM, Swetzes C, et al. Association between health literacy and HIV treatment adherence: further evidence from objectively measured medication adherence. *J Int Assoc Physicians AIDS Care (Chic)* 2008;7:317–23.
- [41] Kripalani S, Gatti ME, Jacobson TA. Association of age, health literacy, and medication management strategies with cardiovascular medication adherence. *Patient Educ Couns* 2010;81:177–81.
- [42] Gazmararian JA, Kripalani S, Miller MJ, Echt KV, Ren J, Rask K. Factors associated with medication refill adherence in cardiovascular-related diseases: a focus on health literacy. *J Gen Intern Med* 2006;21:1215–21.
- [43] Gehi AK, Ali S, Na B, Whooley MA. Self-reported medication adherence and cardiovascular events in patients with stable coronary heart disease: the heart and soul study. *Arch Intern Med* 2007;167:1798–803.
- [44] Schroeder K, Fahey T, Hay AD, Montgomery A, Peters TJ. Adherence to antihypertensive medication assessed by self-report was associated with electronic monitoring compliance. *J Clin Epidemiol* 2006;59:650–1.
- [45] Reyna VF, Nelson WL, Han PK, Dieckmann NF. How numeracy influences risk comprehension and medical decision making. *Psychol Bull* 2009;135:943–73.
- [46] Cavanaugh K, Wallston KA, Gebretsadik T, Shintani A, Huizinga MM, Davis D, et al. Addressing literacy and numeracy to improve diabetes care: two randomized controlled trials. *Diabetes Care* 2009;32:2149–55.
- [47] Cavanaugh K, Davis D, Gregory RP, Wolff K, Rothman RL. Addressing the role of nutrition education and health literacy in diabetes care. In: *American Association of Diabetes Educators Annual Meeting*; 2010.